

MARSHALL, GERSTEIN & BORUN

ATTORNEYS AT LAW

6300 SEARS TOWER

233 SOUTH WACKER DRIVE

CHICAGO, ILLINOIS 60606-6357

(312) 474-6300

FAX: (312) 474-0448

February 17, 2003

FACSIMILE TRANSMISSION SHEET

TO Examiner Hope
Robinson
COMPANY U.S. Patent &
Trademark Office
FAX NO. 703/746-5128
PHONE NO.

FROM: Mark Hopkins, Ph.D.

EXTENSION: 637

PAGES (INCLUDING THIS PAGE): 6

CLIENT NO: 28594

MATTER NO: 35007A

PLEASE CONFIRM RECEIPT: YES

COUNTRY CODE: US

MESSAGE: For U.S. Patent Application Serial No. 09/400,769: **INFORMAL SUBMISSION—
NOT FOR ENTRY INTO FILE**

Please contact Terre Robertson at (312) 474-6825 if you do not receive all of the pages in good condition.

The material of this transmission contains confidential information intended only for the addressee. If you are not the addressee, any disclosure or use of this information by you is strictly prohibited. If you have received this facsimile in error, please notify us by telephone immediately.

INFORMAL SUBMISSION NOT FOR ENTRY INTO FILE

To: Examiner Hope Robinson
 From: Mark Hopkins
 Date: February 14, 2003
 Re: USSN 09/400,769

AMENDMENTSIn the Claims:

Claims 2, 4 and 20-31 will be amended as shown below:

2. A method according to claim 1, wherein the ionic amino acid residue is selected from the group consisting of: A21 Asn, B21 Glu and A17 Glu.
4. A method according to claim 1, wherein at least one amino acid is selected from the group consisting of: A17 Glu, B21 Glu and A21 Asn; and at least one amino acid is selected from the group consisting of: B24 Phe, B25 Phe, A19 Tyr, B12 Val and B12 Tyr.
20. A method according to claim 6 wherein V_1 is selected from the group consisting of: benzene, pyridine, pyridazine, pyrimidine, pyrazine, triazine.
21. A method according to claim 6 wherein V_2 is selected from the group consisting of: cyclopenta-1,3-diene, pyrrole, furan, thiophene, oxazole, isoxazole, pyrazole, imidazole, thiazole, isothiazole or triazole, optionally substituted with up to 4 R_1 groups.
22. A method according to claim 6 wherein W_2 is selected from the group consisting of: naphthalene, quinoline, isoquinoline, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline, pteridine, indole, benzothiophene, benzofuran, benzimidazole, indazole, benzoxazole, benzisooxazole, benzthiazole, benzisothiazole, purine, indoline, isoindoline.
23. A method according to claim 6 wherein R_2 and R'_2 are joined to form cyclic structures selected from the group consisting of: pyrrolidine, piperidine, hexahydro-1H-azepine, morpholine or piperazine.
24. A method according to claim 6 wherein Y_1 is selected from the group consisting of: croman, isochroman, benzofuran, cromene, 1,2,3,4-tetrahydronaphthalene, 1,4-dihydronaphthalene, indan, indene, benzopiperidine, indoline, isoindoline, quinoline,